

## 14.4: Directional Derivatives and the Gradient Vector

### Calculus III

College of the Atlantic. Winter 2016

1. Let  $f(x, y) = 3x^3y^2$ .

(a) Find the gradient of  $f$  at  $x = 1, y = 2$ .

(b) Find the directional derivative  $f_{\vec{u}}$  at  $(1, 2)$  for the following  $\vec{u}$ 's:

i.  $\vec{u} = \vec{i}$

ii.  $\vec{u} = \vec{j}$

iii.  $\vec{u} = \vec{i} + \vec{j}$

iv.  $\vec{u} = \vec{i} - \vec{j}$

v.  $\vec{u} = -\vec{i}$

vi.  $\vec{u} = -\vec{j}$

(c) In what direction  $\vec{u}$  is  $f_{\vec{u}}$  the largest?

(d) In what direction  $\vec{u}$  is  $f_{\vec{u}}$  the smallest?

(e) In what direction  $\vec{u}$  is  $f_{\vec{u}}$  the zero?

2. Consider the function  $f(x, y) = x^2 + 4y^2$ .

(a) Sketch contour lines for the function in the first quadrant.

(b) Calculate the gradient vector for general  $x, y$ .

(c) Determine the value of the gradient vector at the following points:

i.  $(1, 1)$

ii.  $(1, 2)$

iii.  $(2, 1)$

iv.  $(2, 2)$

(d) Draw the above gradient vectors on your contour plot sketch. Do the values make sense geometrically?

(e) What is the rate of change of  $f$  at  $(2, 2)$  in the direction  $\vec{u} = -\vec{i} + 2\vec{j}$ ?

(f) In what direction is the rate of change of  $f$  at  $(2, 2)$  the largest? I.e., in what direction is the function the steepest uphill?

(g) In what direction is the rate of change of  $f$  at  $(2, 2)$  the smallest? I.e., in what direction is the function the steepest downhill?

(h) In what direction is the rate of change of  $f$  at  $(2, 2)$  zero? I.e., in what direction does the function not change?

3. A caterpillar is on a metal surface whose temperature is given by  $T(x, y) = 3x^2y - y^3$ . The caterpillar does not like heat. It is at the point  $(5, 1)$ .
- (a) In what direction should it move so that it gets cooler as quickly as possible?
  - (b) If it initially moves at 0.8 cm/s, at what rate does the caterpillar experience a temperature decrease?
4. A bird is flying through a large cloud of pollution whose distribution is given by  $\rho(x, y, z) = xz + 3x^2y - y^3$  in units of grams per cubic meter, where  $x$ ,  $y$ , and  $z$  are measured in miles. The bird does not like pollution. It is at the point  $(1, 2, 1)$ .
- (a) In what direction should it move so that it gets to cleaner air as quickly as possible?
  - (b) What are the units of the gradient vector?
  - (c) If it initially flies at 1.2 m/s, at what rate does the bird experience a pollution decrease?
5. Consider the function  $f(x, y, z) = e^{-(x^2+y^2+z^2)}$ .
- (a) Calculate  $\vec{\nabla}f$ .
  - (b) Evaluate the gradient function at the points  $(1, 0, 0)$ ,  $(1, 1, 1)$ , and  $(1, -1, 1)$ .
  - (c) What is the gradient vector at the origin? What does your answer mean?