

20.3 and 20.4: Curl and Stokes' Theorem

Calculus III

College of the Atlantic. Winter 2016

- Let $\vec{G} = 2y\hat{i} - 2x\hat{j}$
 - Sketch or describe the field.
 - Calculate $\nabla \times \vec{G}$ and make a sketch of it.
 - Use Stokes' Theorem to calculate $\int_C \vec{G} \cdot d\vec{r}$ where:
 - C is a circle parallel to the yz -plane of radius a , centered at the origin, oriented counter clockwise when viewed from the positive x axis.
 - C is a circle parallel to the xy -plane of radius a , centered at a point on the z -axis, oriented counter clockwise when viewed from above.

- Use Stokes's theorem to find the circulation of \vec{F} around the circle $x^2 + y^2 = 4$, $z = 1$, oriented counterclockwise when viewed from above, and where

$$\vec{F} = (z - 2y)\hat{i} + (3x - 4y)\hat{j} + (z + 3y)\hat{k}. \quad (1)$$

- Use the divergence theorem to find the flux of \vec{F} , given above in Eq. (1), out of a sphere of radius 3 centered at the origin.
- Let $\vec{F}(x, y, z)$ be a vector field and $f(x, y, z)$ is a scalar function of three variables. Which of the following quantities are vectors and which are scalars. Which are not defined?

- $\text{div } \vec{F}$
- $\text{curl } \vec{F}$
- $\text{div } f$
- $\text{curl } f$
- ∇f
- $\nabla \vec{F}$
- $\nabla \times \vec{F}$
- $\nabla \cdot \vec{F}$
- $\nabla \cdot f$
- $\nabla \times f$
- $\nabla \cdot \nabla f$
- $\nabla \times \nabla f$
- $\nabla \cdot \nabla \times \vec{F}$