

## C8: Force and Energy

This chapter is about *relationships...*

### C8.1: Momentum and Kinetic Energy

Kinetic energy is related to momentum:

$$K = \frac{p^2}{2m} . \quad (1)$$

Not all momentum transfers lead to a change in kinetic energy. Consider a small momentum transfer  $d\vec{p}$ . Suppose an object is moving at velocity  $\vec{v}$ . Only the “portion” of  $d\vec{p}$  that is in the same direction as  $\vec{v}$  will lead to a change in kinetic energy.

$$dK = v dp \cos \theta = \vec{v} \cdot d\vec{p} , \quad (2)$$

where  $v$  is the speed of the object,  $dp$  the magnitude of the momentum transfer, and  $\theta$  is the angle between  $\vec{v}$  and  $d\vec{p}$ .

### C8.2: The Dot Product

The dot product between  $\vec{u}$  and  $\vec{v}$  is the magnitude of  $\vec{u}$  times that portion of  $\text{mag}(\vec{v})$  that’s in  $\vec{u}$ ’s direction.

Two important formulas:

$$\vec{u} \cdot \vec{w} \equiv uv \cos \theta , \quad (3)$$

where  $\theta \equiv$  the angle between  $\vec{u}$  and  $\vec{v}$ .

Also,

$$\vec{u} \cdot \vec{w} = u_x w_x + u_y w_y + u_z w_z . \quad (4)$$

Note that:

- $\vec{u} \cdot \vec{w}$  is a scalar.
- $\vec{u} \cdot \vec{w}$  can be positive or negative.
- $\vec{u} \cdot \vec{u} = u^2$ .

### **C8.3 An Interaction's Contribution to $dK$**

An interaction gives rise to a force on an object. The amount by which this interaction changes the object's kinetic energy is given by:

$$dK \equiv \vec{F} \cdot d\vec{r} \quad (5)$$

### **C8.4 The Meaning of k-Work**

When there's a kinetic energy transfer  $dK$ , the energy comes from some sort of potential energy – it does not come from another interaction. Remember that energy is a property of an interaction, not a property of a particular object.

### **C8.5 The Earth's Kinetic Energy**

Yet again, we note that the earth is way bigger than us.

### **C8.6 Force Laws**

Don't worry about this section. The main point is that one can go from a potential energy function to a force and vice-versa.

### **C8.7 Contact Interactions**

The normal (perpendicular) part of a contact interaction contributes no k-work.

## Practice

1. Consider two displacement vectors:  $\vec{v}_1 = [2m, -4m]$  and  $\vec{v}_2 = [3m, -1m]$ . Calculate  $\vec{v}_1 \cdot \vec{v}_2$ . Calculate the angle between  $\vec{v}_1$  and  $\vec{v}_2$ .
2. A 5 kg object is traveling due north at 10 m/s.
  - (a) The object is briefly acted upon by a force of 2 Newtons due east. This force displaces the object 2 cm. What energy transfer has the object received?
  - (b) The object is briefly acted upon by a force of 2 Newtons due south. This force displaces the object 2 cm. What energy transfer has the object received?
  - (c) The object is briefly acted upon by a force of 2 Newtons 37 degrees west of north. This force displaces the object 2 cm. What energy transfer has the object received?
3. A 2000 kg car rolls down a 37 degree incline at a constant speed of 20 m/s.
  - (a) In one second, what energy transfer does the gravitational interaction give to the car?
  - (b) Where does this energy transfer go?
4. A car goes over the crest of a hill at 20 m/s. The car then coasts to the bottom of the hill, 50 meters below. Ignoring friction, what is the car's speed at the bottom?